



Generalized fractional midpoint type inequalities for co-ordinated convex functions

Fatih Hezenci^a, Hüseyin Budak^a, Hasan Kara^a, Mehmet Zeki Sarıkaya^a

^aDepartment of Mathematics, Faculty of Science and Arts, Duzce University, Türkiye

Abstract. In this research paper, we investigate generalized fractional integrals to obtain midpoint type inequalities for the co-ordinated convex functions. First of all, we establish an identity for twice partially differentiable mappings. By utilizing this equality, some midpoint type inequalities via generalized fractional integrals are proved. We also show that the main results reduce some midpoint inequalities given in earlier works for Riemann integrals and Riemann-Liouville fractional integrals. Finally, some new inequalities for k -Riemann-Liouville fractional integrals are presented as special cases of our results.

1. Introduction

The inequalities, introduced by C. Hermite and J. Hadamard for convex functions, are significant issue in the literature. These inequalities state that if $f : I \rightarrow \mathbb{R}$ is a convex function on the interval I of real numbers and $a, b \in I$ with $a < b$, then the following double inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2} \quad (1)$$

is valid. If f is concave, then both inequalities in (1) hold to the reverse direction. With the help of the convex functions, Dragomir and Agarwal [12] first obtained an upper bound for

$$\frac{f(a)+f(b)}{2} - \frac{1}{b-a} \int_a^b f(x)dx,$$

which is the right-hand side of inequality (1). In addition to this, Kirmacı [20] first obtained upper bound for

$$\frac{1}{b-a} \int_a^b f(x)dx - f\left(\frac{a+b}{2}\right),$$

2020 Mathematics Subject Classification. Primary 26D07, 26D10, 26D15; Secondary 26B15, 26B25.

Keywords. Simpson type inequalities, Convex function, Fractional integrals, Third derivative

Received: 13 July 2022; Revised: 31 August 2022; Accepted: 12 October 2022

Communicated by Eberhard Malkowsky

Email addresses: fatihezenci@gmail.com (Fatih Hezenci), hsyn.budak@gmail.com (Hüseyin Budak), [\(Hasan Kara\)](mailto:hasan64kara@gmail.com), sarikayamz@gmail.com (Mehmet Zeki Sarıkaya)

which is the left-hand side of inequality (1). These inequalities are called by trapezoid type inequality and midpoint type inequality, respectively. Many researchers have been studied extensively the trapezoid inequalities and midpoint inequalities for various types of convex functions [3, 21, 28, 33, 35, 45]. In 2013, Sarikaya et al. first proved Hermite-Hadamard inequalities for Riemann-Liouville fractional integrals and the authors also gave some corresponding trapezoid type inequalities [39]. With the help of the results of Sarikaya et al., some fractional midpoint type inequalities for convex functions were established in [17]. Thereupon, many researchers obtained fractional midpoint inequalities and trapezoid inequalities using different kind of fractional integrals [1, 2, 4, 5, 8, 11, 27, 29, 32, 36, 37, 46]. On the other hand, Dragomir first proved Hermite-Hadamard inequalities for co-ordinated convex mappings in [13]. The midpoint and trapezoid type inequalities for co-ordinated convex functions were established in the papers [23] and [38], respectively. Furthermore, the Hermite-Hadamard inequalities for functions with two variables by utilizing Riemann-Liouville fractional integrals were obtained in [40]. Although Sarikaya gave the corresponding fractional trapezoid inequalities for co-ordinated convex functions in [40], Tunç et al. presented fractional midpoint type inequalities for co-ordinated convex functions in [43]. In the literature, there are great number of papers about to Hermite-Hadamard inequalities for several type co-ordinated convex functions. Because of these reasons, the reader is referred to [9, 10, 19, 22, 24, 30, 31, 42] and the references therein for additional information and unexplained subjects about these topics.

The generalized fractional integrals, which will be used frequently throughout this paper, were introduced by Sarikaya and Ertuğral in [41]. In the same paper, the authors also established Hermite-Hadamard inequalities and introduced several trapezoid and midpoint type inequalities for this kind fractional integrals. In addition to this, Turkay et al. defined the generalized fractional integrals for functions with two variables in [44]. These authors presented Hermite-Hadamard and trapezoid type inequalities for this kind of fractional integrals.

The purpose of this paper is to establish some generalized midpoint type inequalities for co-ordinated convex functions involving generalized fractional integrals. The general structure of our article contains four parts, including the introduction. The rest of the paper continues as follows: In Section 2, the definitions of generalized fractional integrals are given. Moreover, relations between generalized fractional integrals and other type fractional integrals are introduced. In Section 3, we first prove an equality involving for twice partially differentiable functions. Then, we establish several generalized midpoint type inequalities whose partially derivatives in absolute value are co-ordinated convex. We show that our main results are reduced to inequalities from earlier studies by looking at their particular case. Furthermore, some new midpoint type inequalities for k -Riemann-Liouville fractional integrals are given. Finally, some results and further aspects of research are discussed in the last section.

2. Generalized fractional Integrals

In this section, we will give the necessary definition of generalized fractional integrals introduced by Sarikaya and Ertuğral in [41]. Moreover, the relation between generalized fractional integrals and other type integrals are considered.

Definition 2.1. Let $f : [a, b] \rightarrow \mathbb{R}$ denote a integrable function. The left-sided and right-sided generalized fractional integral operators are given by

$${}_{a+}I_\varphi f(x) = \int_a^x \frac{\varphi(x-t)}{x-t} f(t) dt, \quad x > a \tag{2}$$

and

$${}_{b-}I_\varphi f(x) = \int_x^b \frac{\varphi(t-x)}{t-x} f(t) dt, \quad x < b, \tag{3}$$

respectively. Here, the function $\varphi : [0, \infty) \rightarrow [0, \infty)$ satisfying the condition

$$\int_0^1 \frac{\varphi(t)}{t} dt < \infty.$$

Remark 2.2. With the help of the given Definition 2.1, the following cases are provided:

1. If we choose $\varphi(t) = t$, the operators (2) and (3) reduce to the Riemann integral.
2. Let us consider $\varphi(t) = \frac{t^\alpha}{\Gamma(\alpha)}$ and $\alpha > 0$. Then, the operators (2) and (3) reduce to the Riemann-Liouville fractional integrals $J_{a+}^\alpha f(x)$ and $J_{b-}^\alpha f(x)$, respectively. Here, Γ is Gamma function.
3. Let us define $\varphi(t) = \frac{1}{k\Gamma_k(\alpha)} t^{\frac{\alpha}{k}}$ and $\alpha, k > 0$. Then, the operators (2) and (3) reduce to the k -Riemann-Liouville fractional integrals $J_{a+k}^\alpha f(x)$ and $J_{b-k}^\alpha f(x)$, respectively. Here, Γ_k is k -Gamma function.

There are several papers on inequalities for generalized fractional integrals in the literature. In [41], Sarikaya and Ertuğral also proved Hermite-Hadamard inequalities for generalized fractional integrals. In addition, Budak et al. proved midpoint type inequalities and extensions of Hermite-Hadamard inequalities in the papers [6] and [7], respectively. In [14], Ertuğral and Sarikaya presented some Simpson type inequalities for these fractional integral operators. For some of other papers on inequalities for generalized fractional integrals, please refer to [15, 16, 18, 25, 26, 34, 47].

Generalized double fractional integrals are given by Turkay et al. in [44], as follows:

Definition 2.3. Let $f : \Omega := [a, b] \times [c, d] \rightarrow \mathbb{R}$ be a integrable function. The generalized double fractional integrals ${}_{a+,c+}I_{\varphi,\psi}$, ${}_{a+,d-}I_{\varphi,\psi}$, ${}_{b-,c+}I_{\varphi,\psi}$, ${}_{b-,d-}I_{\varphi,\psi}$ are defined by

$${}_{a+,c+}I_{\varphi,\psi} f(x, y) = \int_a^x \int_c^y \frac{\varphi(x-t)}{x-t} \frac{\psi(y-s)}{y-s} f(t, s) ds dt, \quad x > a, y > c, \quad (4)$$

$${}_{a+,d-}I_{\varphi,\psi} f(x, y) = \int_a^x \int_y^d \frac{\varphi(x-t)}{x-t} \frac{\psi(s-y)}{s-y} f(t, s) ds dt, \quad x > a, y < d, \quad (5)$$

$${}_{b-,c+}I_{\varphi,\psi} f(x, y) = \int_x^b \int_c^y \frac{\varphi(t-x)}{t-x} \frac{\psi(y-s)}{y-s} f(t, s) ds dt, \quad x < b, y > c, \quad (6)$$

and

$${}_{b-,d-}I_{\varphi,\psi} f(x, y) = \int_x^b \int_y^d \frac{\varphi(t-x)}{t-x} \frac{\psi(s-y)}{s-y} f(t, s) ds dt, \quad x < b, y < d. \quad (7)$$

Here, the function $\varphi : [0, \infty) \rightarrow [0, \infty)$ and the function $\psi : [0, \infty) \rightarrow [0, \infty)$ satisfy the conditions $\int_0^1 \frac{\varphi(t)}{t} dt < \infty$ and $\int_0^1 \frac{\psi(s)}{s} ds < \infty$, respectively.

Remark 2.4. By using the Definition 2.3, the following conditions are ensured:

1. If we take $\varphi(t) = t$ and $\psi(s) = s$, then the operators (4), (5), (6), and (7) reduce to the double Riemann integral.
2. Let us consider $\varphi(t) = \frac{t^\alpha}{\Gamma(\alpha)}$, $\psi(s) = \frac{s^\beta}{\Gamma(\beta)}$ for $\alpha, \beta > 0$. Then, the operators (4), (5), (6), and (7) reduce to the Riemann-Liouville fractional integrals $J_{a+,c+}^{\alpha,\beta} f(x, y)$, $J_{a+,d-}^{\alpha,\beta} f(x, y)$, $J_{b-,c+}^{\alpha,\beta} f(x, y)$, and $J_{b-,d-}^{\alpha,\beta} f(x, y)$, respectively.
3. Let us note that $\varphi(t) = \frac{t^{\frac{\alpha}{k}}}{k\Gamma_k(\alpha)}$ and $\psi(s) = \frac{s^{\frac{\beta}{k}}}{k\Gamma_k(\beta)}$ for $\alpha, \beta, k > 0$. Then, the operators (4), (5), (6), and (7) reduce to the k -Riemann-Liouville fractional integrals $J_{a+,c+}^{\alpha,\beta,k} f(x, y)$, $J_{a+,d-}^{\alpha,\beta,k} f(x, y)$, $J_{b-,c+}^{\alpha,\beta,k} f(x, y)$, and $J_{b-,d-}^{\alpha,\beta,k} f(x, y)$, respectively.

3. Generalized midpoint type Inequalities for co-ordinated convex functions

To make the presentation easier and compact to understand, we make some symbolic representation:

$$\begin{aligned}
 I_1 &= \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} \Lambda_1(t) \Lambda_2(s) \frac{\partial^2 f}{\partial t \partial s} (ta + (1-t)b, sc + (1-s)d) ds dt, \\
 I_2 &= - \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 \Lambda_1(t) \Delta_2(s) \frac{\partial^2 f}{\partial t \partial s} (ta + (1-t)b, sc + (1-s)d) ds dt, \\
 I_3 &= - \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} \Delta_1(t) \Lambda_2(s) \frac{\partial^2 f}{\partial t \partial s} (ta + (1-t)b, sc + (1-s)d) ds dt, \\
 I_4 &= \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 \Delta_1(t) \Delta_2(s) \frac{\partial^2 f}{\partial t \partial s} (ta + (1-t)b, sc + (1-s)d) ds dt, \\
 I_5 &= \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} \Lambda_1(t) \Lambda_2(s) \frac{\partial^2 f}{\partial t \partial s} (tb + (1-t)a, sd + (1-s)c) ds dt, \\
 I_6 &= - \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 \Lambda_1(t) \Delta_2(s) \frac{\partial^2 f}{\partial t \partial s} (tb + (1-t)a, sd + (1-s)c) ds dt, \\
 I_7 &= - \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} \Delta_1(t) \Lambda_2(s) \frac{\partial^2 f}{\partial t \partial s} (tb + (1-t)a, sd + (1-s)c) ds dt, \\
 I_8 &= \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 \Delta_1(t) \Delta_2(s) \frac{\partial^2 f}{\partial t \partial s} (tb + (1-t)a, sd + (1-s)c) ds dt, \\
 I_9 &= - \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} \Lambda_1(t) \Lambda_2(s) \frac{\partial^2 f}{\partial t \partial s} (ta + (1-t)b, sd + (1-s)c) ds dt, \\
 I_{10} &= \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 \Lambda_1(t) \Delta_2(s) \frac{\partial^2 f}{\partial t \partial s} (ta + (1-t)b, sd + (1-s)c) ds dt, \\
 I_{11} &= \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} \Delta_1(t) \Lambda_2(s) \frac{\partial^2 f}{\partial t \partial s} (ta + (1-t)b, sd + (1-s)c) ds dt, \\
 I_{12} &= - \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 \Delta_1(t) \Delta_2(s) \frac{\partial^2 f}{\partial t \partial s} (ta + (1-t)b, sd + (1-s)c) ds dt, \\
 I_{13} &= - \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} \Lambda_1(t) \Lambda_2(s) \frac{\partial^2 f}{\partial t \partial s} (tb + (1-t)a, sc + (1-s)d) ds dt, \\
 I_{14} &= \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 \Lambda_1(t) \Delta_2(s) \frac{\partial^2 f}{\partial t \partial s} (tb + (1-t)a, sc + (1-s)d) ds dt, \\
 I_{15} &= \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} \Delta_1(t) \Lambda_2(s) \frac{\partial^2 f}{\partial t \partial s} (tb + (1-t)a, sc + (1-s)d) ds dt, \\
 I_{16} &= - \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 \Delta_1(t) \Delta_2(s) \frac{\partial^2 f}{\partial t \partial s} (tb + (1-t)a, sc + (1-s)d) ds dt,
 \end{aligned}$$

where

$$\begin{cases} \Lambda_1(t) = \int_0^t \frac{\varphi((b-a)u)}{u} du, \\ \Lambda_2(s) = \int_0^s \frac{\psi((d-c)u)}{u} du, \end{cases}$$

and

$$\begin{cases} \Delta_1(t) = \int_t^1 \frac{\varphi((b-a)u)}{u} du, \\ \Delta_2(s) = \int_s^1 \frac{\psi((d-c)u)}{u} du. \end{cases}$$

In order to prove our main results, we need the following Lemma.

Lemma 3.1. *Let $f : \Omega \rightarrow \mathbb{R}$ be a partial differentiable mapping on Ω and let $\frac{\partial^2 f}{\partial t \partial s} \in L(\Omega)$. Then, the following equality holds:*

$$\begin{aligned} & f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{1}{2\Lambda_1(1)} \left[{}_{a+}I_\varphi f\left(b, \frac{c+d}{2}\right) + {}_{b-}I_\varphi f\left(a, \frac{c+d}{2}\right) \right] \\ & - \frac{1}{2\Lambda_2(1)} \left[{}_{c+}I_\psi f\left(\frac{a+b}{2}, d\right) + {}_{d-}I_\psi f\left(\frac{a+b}{2}, c\right) \right] \\ & + \frac{1}{4\Lambda_1(1)\Lambda_2(1)} \left[{}_{a+,c+}I_{\varphi,\psi} f(b, d) + {}_{a+,d-}I_{\varphi,\psi} f(b, c) + {}_{b-,c+}I_{\varphi,\psi} f(a, d) + {}_{b-,d-}I_{\varphi,\psi} f(a, c) \right] \\ & = \frac{(b-a)(d-c)}{4\Lambda_1(1)\Lambda_2(1)} \sum_{k=1}^{16} I_k. \end{aligned} \quad (8)$$

Proof. With the help of the integration by parts, we obtain

$$\begin{aligned} I_1 &= \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} \Lambda_1(t) \Lambda_2(s) \frac{\partial^2 f}{\partial t \partial s}(ta + (1-t)b, sc + (1-s)d) ds dt \\ &= \int_0^{\frac{1}{2}} \Lambda_1(t) \left\{ \Lambda_2(s) \frac{1}{c-d} \frac{\partial f}{\partial t}(ta + (1-t)b, sc + (1-s)d) \Big|_0^{\frac{1}{2}} \right. \\ &\quad \left. - \frac{1}{c-d} \int_0^{\frac{1}{2}} \frac{\psi((d-c)s)}{s} \frac{\partial f}{\partial t}(ta + (1-t)b, sc + (1-s)d) ds \right\} dt \\ &= \frac{\Lambda_2\left(\frac{1}{2}\right)}{c-d} \int_0^{\frac{1}{2}} \Lambda_1(t) \frac{\partial f}{\partial t}\left(ta + (1-t)b, \frac{c+d}{2}\right) dt \\ &\quad - \frac{1}{c-d} \int_0^{\frac{1}{2}} \frac{\psi((d-c)s)}{s} \int_0^{\frac{1}{2}} \Lambda_1(t) \frac{\partial f}{\partial t}(ta + (1-t)b, sc + (1-s)d) dt ds \\ &= \frac{\Lambda_2\left(\frac{1}{2}\right)}{c-d} \left[\frac{\Lambda_1\left(\frac{1}{2}\right)}{a-b} f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{1}{a-b} \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} f\left(ta + (1-t)b, \frac{c+d}{2}\right) dt \right] \end{aligned} \quad (9)$$

$$\begin{aligned}
& -\frac{1}{c-d} \int_0^{\frac{1}{2}} \frac{\psi((d-c)s)}{s} \left[\frac{\Lambda_1\left(\frac{1}{2}\right)}{a-b} f\left(\frac{a+b}{2}, sc + (1-s)d\right) \right. \\
& \quad \left. - \frac{1}{a-b} \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} \frac{\partial f}{\partial s}(ta + (1-t)b, sc + (1-s)d) dt \right] ds \\
& = \frac{1}{(b-a)(d-c)} \left[\Lambda_1\left(\frac{1}{2}\right) \Lambda_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\
& \quad \left. - \Lambda_2\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} f\left(ta + (1-t)b, \frac{c+d}{2}\right) dt \right. \\
& \quad \left. - \Lambda_1\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sc + (1-s)d\right) ds \right. \\
& \quad \left. + \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(ta + (1-t)b, sc + (1-s)d) ds dt \right].
\end{aligned}$$

Similarly, by using integration by parts it follows that

$$\begin{aligned}
I_2 &= \frac{1}{(b-a)(d-c)} \left[\Lambda_1\left(\frac{1}{2}\right) \Delta_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\
&\quad \left. - \Delta_2\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} f\left(ta + (1-t)b, \frac{c+d}{2}\right) dt \right. \\
&\quad \left. - \Lambda_1\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sc + (1-s)d\right) ds \right. \\
&\quad \left. + \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(ta + (1-t)b, sc + (1-s)d) ds dt \right], \tag{10}
\end{aligned}$$

$$\begin{aligned}
I_3 &= \frac{1}{(b-a)(d-c)} \left[\Delta_1\left(\frac{1}{2}\right) \Lambda_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\
&\quad \left. - \Lambda_2\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} f\left(ta + (1-t)b, \frac{c+d}{2}\right) dt \right. \\
&\quad \left. - \Delta_1\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sc + (1-s)d\right) ds \right. \\
&\quad \left. + \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(ta + (1-t)b, sc + (1-s)d) ds dt \right], \tag{11}
\end{aligned}$$

$$\begin{aligned}
I_4 &= \frac{1}{(b-a)(d-c)} \left[\Delta_1\left(\frac{1}{2}\right) \Delta_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\
&\quad \left. - \Delta_2\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} f\left(ta + (1-t)b, \frac{c+d}{2}\right) dt \right] \tag{12}
\end{aligned}$$

$$\begin{aligned} & -\Delta_1\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sc + (1-s)d\right) ds \\ & + \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(ta + (1-t)b, sc + (1-s)d) ds dt \Big], \end{aligned}$$

$$\begin{aligned} I_5 = & \frac{1}{(b-a)(d-c)} \left[\Lambda_1\left(\frac{1}{2}\right) \Lambda_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ & -\Lambda_2\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} f\left(tb + (1-t)a, \frac{c+d}{2}\right) dt \\ & -\Lambda_1\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sd + (1-s)c\right) ds \\ & \left. + \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(tb + (1-t)a, sd + (1-s)c) ds dt \right], \end{aligned} \quad (13)$$

$$\begin{aligned} I_6 = & \frac{1}{(b-a)(d-c)} \left[\Lambda_1\left(\frac{1}{2}\right) \Delta_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ & -\Delta_2\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} f\left(tb + (1-t)a, \frac{c+d}{2}\right) dt \\ & -\Lambda_1\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sd + (1-s)c\right) ds \\ & \left. + \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(tb + (1-t)a, sd + (1-s)c) ds dt \right], \end{aligned} \quad (14)$$

$$\begin{aligned} I_7 = & \frac{1}{(b-a)(d-c)} \left[\Delta_1\left(\frac{1}{2}\right) \Lambda_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ & -\Lambda_2\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} f\left(tb + (1-t)a, \frac{c+d}{2}\right) dt \\ & -\Delta_1\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sd + (1-s)c\right) ds \\ & \left. + \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(tb + (1-t)a, sd + (1-s)c) ds dt \right], \end{aligned} \quad (15)$$

$$I_8 = \frac{1}{(b-a)(d-c)} \left[\Delta_1\left(\frac{1}{2}\right) \Delta_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \quad (16)$$

$$\begin{aligned} & -\Delta_2\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} f\left(tb + (1-t)a, \frac{c+d}{2}\right) dt \\ & -\Delta_1\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sd + (1-s)c\right) ds \\ & \left. + \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(tb + (1-t)a, sd + (1-s)c) ds dt \right], \end{aligned} \quad (17)$$

$$I_9 = \frac{1}{(b-a)(d-c)} \left[\Lambda_1\left(\frac{1}{2}\right) \Lambda_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ - \Lambda_2\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} f\left(ta + (1-t)b, \frac{c+d}{2}\right) dt \\ - \Lambda_1\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sd + (1-s)c\right) ds \\ \left. + \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(ta + (1-t)b, sd + (1-s)c) ds dt \right], \quad (18)$$

$$I_{10} = \frac{1}{(b-a)(d-c)} \left[\Lambda_1\left(\frac{1}{2}\right) \Lambda_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ - \Lambda_2\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} f\left(ta + (1-t)b, \frac{c+d}{2}\right) dt \\ - \Lambda_1\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sd + (1-s)c\right) ds \\ \left. + \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(ta + (1-t)b, sd + (1-s)c) ds dt \right], \quad (19)$$

$$I_{11} = \frac{1}{(b-a)(d-c)} \left[\Lambda_1\left(\frac{1}{2}\right) \Lambda_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ - \Lambda_2\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} f\left(ta + (1-t)b, \frac{c+d}{2}\right) dt \\ - \Lambda_1\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sd + (1-s)c\right) ds \\ \left. + \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(ta + (1-t)b, sd + (1-s)c) ds dt \right], \quad (20)$$

$$I_{12} = \frac{1}{(b-a)(d-c)} \left[\Lambda_1\left(\frac{1}{2}\right) \Lambda_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ - \Lambda_2\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} f\left(ta + (1-t)b, \frac{c+d}{2}\right) dt \\ - \Lambda_1\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sd + (1-s)c\right) ds \\ \left. + \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(ta + (1-t)b, sd + (1-s)c) ds dt \right], \quad (21)$$

$$I_{13} = \frac{1}{(b-a)(d-c)} \left[\Lambda_1\left(\frac{1}{2}\right) \Lambda_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ - \Lambda_2\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} f\left(tb + (1-t)a, \frac{c+d}{2}\right) dt \quad (22)$$

$$\begin{aligned} & -\Lambda_1\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sc + (1-s)d\right) ds \\ & + \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(tb + (1-t)a, sc + (1-s)d) ds dt \Big], \end{aligned}$$

$$\begin{aligned} I_{14} = & \frac{1}{(b-a)(d-c)} \left[\Lambda_1\left(\frac{1}{2}\right) \Delta_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ & - \Delta_2\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} f\left(tb + (1-t)a, \frac{c+d}{2}\right) dt \\ & - \Lambda_1\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sc + (1-s)d\right) ds \\ & \left. + \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(tb + (1-t)a, sc + (1-s)d) ds dt \right], \end{aligned} \quad (23)$$

$$\begin{aligned} I_{15} = & \frac{1}{(b-a)(d-c)} \left[\Delta_1\left(\frac{1}{2}\right) \Lambda_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ & - \Lambda_2\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} f\left(tb + (1-t)a, \frac{c+d}{2}\right) dt \\ & - \Delta_1\left(\frac{1}{2}\right) \int_0^{\frac{1}{2}} \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sc + (1-s)d\right) ds \\ & \left. + \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(tb + (1-t)a, sc + (1-s)d) ds dt \right], \end{aligned} \quad (24)$$

and

$$\begin{aligned} I_{16} = & \frac{1}{(b-a)(d-c)} \left[\Delta_1\left(\frac{1}{2}\right) \Delta_2\left(\frac{1}{2}\right) f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ & - \Delta_2\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} f\left(tb + (1-t)a, \frac{c+d}{2}\right) dt \\ & - \Delta_1\left(\frac{1}{2}\right) \int_{\frac{1}{2}}^1 \frac{\psi((d-c)s)}{s} f\left(\frac{a+b}{2}, sc + (1-s)d\right) ds \\ & \left. + \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 \frac{\varphi((b-a)t)}{t} \frac{\psi((d-c)s)}{s} f(tb + (1-t)a, sc + (1-s)d) ds dt \right]. \end{aligned} \quad (25)$$

By using the equations (9)-(25), the change of the variable $x = ta + (1-t)b$, and $y = sc + (1-s)d$ for $t, s \in [0, 1]$, it can be rewritten as follows

$$\begin{aligned} & I_1 + I_2 + \cdots + I_{16} \\ & = \frac{4\Lambda_1(1)\Lambda_2(1)}{(b-a)(d-c)} f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \end{aligned} \quad (26)$$

$$\begin{aligned}
& - \frac{2\Lambda_2(1)}{(b-a)(d-c)} \left[{}_{a+}I_\varphi f \left(b, \frac{c+d}{2} \right) + {}_{b-}I_\varphi f \left(a, \frac{c+d}{2} \right) \right] \\
& - \frac{2\Lambda_1(1)}{(b-a)(d-c)} \left[{}_{c+}I_\psi f \left(\frac{a+b}{2}, d \right) + {}_{d-}I_\psi f \left(\frac{a+b}{2}, c \right) \right] \\
& + \frac{1}{(b-a)(d-c)} \left[{}_{a+,c+}I_{\varphi,\psi} f(b, d) + {}_{a+,d-}I_{\varphi,\psi} f(b, c) + {}_{b-,c+}I_{\varphi,\psi} f(a, d) + {}_{b-,d-}I_{\varphi,\psi} f(a, c) \right].
\end{aligned}$$

Multiplying the both sides of (26) by $\frac{(b-a)(d-c)}{4\Lambda_1(1)\Lambda_2(1)}$, we get equation (8). This ends the proof of Lemma 3.1. \square

Next, we start to state the first theorem containing the Hermite-Hadamard type inequality for fractional integrals.

Theorem 3.2. Suppose $f : \Omega \rightarrow \mathbb{R}$ is a partial differentiable mapping on Ω . Suppose also $\left| \frac{\partial^2 f}{\partial t \partial s} \right|$ is a convex function on the co-ordinates on Ω . Then, the following inequality

$$\begin{aligned}
& \left| f \left(\frac{a+b}{2}, \frac{c+d}{2} \right) - \frac{1}{2\Lambda_1(1)} \left[{}_{a+}I_\varphi f \left(b, \frac{c+d}{2} \right) + {}_{b-}I_\varphi f \left(a, \frac{c+d}{2} \right) \right] \right. \\
& - \frac{1}{2\Lambda_2(1)} \left[{}_{c+}I_\psi f \left(\frac{a+b}{2}, d \right) + {}_{d-}I_\psi f \left(\frac{a+b}{2}, c \right) \right] \\
& \left. + \frac{1}{4\Lambda_1(1)\Lambda_2(1)} \left[{}_{a+,c+}I_{\varphi,\psi} f(b, d) + {}_{a+,d-}I_{\varphi,\psi} f(b, c) + {}_{b-,c+}I_{\varphi,\psi} f(a, d) + {}_{b-,d-}I_{\varphi,\psi} f(a, c) \right] \right| \\
& \leq \frac{(b-a)(d-c)}{4\Lambda_1(1)\Lambda_2(1)} [A_1 + A_2][B_1 + B_2] \\
& \times \left(\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right| + \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right| + \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right| + \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right| \right)
\end{aligned}$$

is valid. Here,

$$\begin{cases} A_1 = \int_0^{\frac{1}{2}} |\Delta_1(t)| dt, & B_1 = \int_0^{\frac{1}{2}} |\Delta_2(s)| ds, \\ A_2 = \int_{\frac{1}{2}}^1 |\Delta_1(t)| dt, & B_2 = \int_{\frac{1}{2}}^1 |\Delta_2(s)| ds. \end{cases}$$

Proof. Lemma 3.1 yields the following inequality

$$\begin{aligned}
& \left| f \left(\frac{a+b}{2}, \frac{c+d}{2} \right) - \frac{1}{2\Lambda_1(1)} \left[{}_{a+}I_\varphi f \left(b, \frac{c+d}{2} \right) + {}_{b-}I_\varphi f \left(a, \frac{c+d}{2} \right) \right] \right. \\
& - \frac{1}{2\Lambda_2(1)} \left[{}_{c+}I_\psi f \left(\frac{a+b}{2}, d \right) + {}_{d-}I_\psi f \left(\frac{a+b}{2}, c \right) \right] \\
& \left. + \frac{1}{4\Lambda_1(1)\Lambda_2(1)} \left[{}_{a+,c+}I_{\varphi,\psi} f(b, d) + {}_{a+,d-}I_{\varphi,\psi} f(b, c) + {}_{b-,c+}I_{\varphi,\psi} f(a, d) + {}_{b-,d-}I_{\varphi,\psi} f(a, c) \right] \right| \tag{27}
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{4\Lambda_1(1)\Lambda_2(1)} \left[{}_{a+,c+}I_{\varphi,\psi}f(b,d) + {}_{a+,d-}I_{\varphi,\psi}f(b,c) + {}_{b-,c+}I_{\varphi,\psi}f(a,d) + {}_{b-,d-}I_{\varphi,\psi}f(a,c) \right] \\
& \leq \frac{(b-a)(d-c)}{4\Lambda_1(1)\Lambda_2(1)} \{ |I_1| + |I_2| + \dots + |I_{16}| \}.
\end{aligned}$$

By using co-ordinated convexity of $\left| \frac{\partial^2 f}{\partial t \partial s} \right|$ function on Ω and calculating the integrals in above inequality, the following inequalities hold:

$$|I_1| \leq \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| \left| \frac{\partial^2 f}{\partial t \partial s} (ta + (1-t)b, sc + (1-s)d) \right| ds dt \quad (28)$$

$$\leq \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| \left\{ ts \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| + s(1-t) \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \right.$$

$$\left. + t(1-s) \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| + (1-s)(1-t) \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \right\} ds dt$$

$$= \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| ts ds dt$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| s(1-t) ds dt$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| t(1-s) ds dt$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| (1-s)(1-t) ds dt,$$

$$|I_2| \leq \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Lambda_1(t)| |\Delta_2(s)| ts ds dt \quad (29)$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Lambda_1(t)| |\Delta_2(s)| s(1-t) ds dt$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Lambda_1(t)| |\Delta_2(s)| t(1-s) ds dt$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Lambda_1(t)| |\Delta_2(s)| (1-s)(1-t) ds dt,$$

$$|I_3| \leq \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Lambda_2(s)| tsdsdt \quad (30)$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Lambda_2(s)| s(1-t) dsdt$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Lambda_2(s)| t(1-s) dsdt$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Lambda_2(s)| (1-s)(1-t) dsdt,$$

$$|I_4| \leq \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| tsdsdt \quad (31)$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| s(1-t) dsdt$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| t(1-s) dsdt$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| (1-s)(1-t) dsdt,$$

$$|I_5| \leq \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| tsdsdt \quad (32)$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| s(1-t) dsdt$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| t(1-s) dsdt$$

$$+ \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| (1-s)(1-t) dsdt,$$

$$|I_6| \leq \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Lambda_1(t)| |\Delta_2(s)| tsdsdt \quad (33)$$

$$\begin{aligned}
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| s(1-t) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| t(1-s) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| (1-s)(1-t) ds dt, \\
|I_7| & \leq \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Delta_2(s)| ts ds dt
\end{aligned} \tag{34}$$

$$\begin{aligned}
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Delta_2(s)| s(1-t) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Delta_2(s)| t(1-s) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Delta_2(s)| (1-s)(1-t) ds dt, \\
|I_8| & \leq \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| ts ds dt
\end{aligned} \tag{35}$$

$$\begin{aligned}
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| s(1-t) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| t(1-s) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| (1-s)(1-t) ds dt, \\
|I_9| & \leq \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Delta_2(s)| ts ds dt
\end{aligned} \tag{36}$$

$$\begin{aligned}
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| t(1-s) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| (1-s)(1-t) ds dt, \\
|I_{10}| & \leq \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_0^{\frac{1}{2}} \int_0^1 |\Lambda_1(t)| |\Delta_2(s)| ts ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Lambda_1(t)| |\Delta_2(s)| s(1-t) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Lambda_1(t)| |\Delta_2(s)| t(1-s) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Lambda_1(t)| |\Delta_2(s)| (1-s)(1-t) ds dt,
\end{aligned} \tag{37}$$

$$\begin{aligned}
|I_{11}| & \leq \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Lambda_2(s)| ts ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Lambda_2(s)| s(1-t) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Lambda_2(s)| t(1-s) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Lambda_2(s)| (1-s)(1-t) ds dt, \\
|I_{12}| & \leq \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| ts ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| s(1-t) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| t(1-s) ds dt
\end{aligned} \tag{38}$$

$$\begin{aligned}
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| (1-s)(1-t) ds dt, \\
|I_{13}| & \leq \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| ts ds dt
\end{aligned} \tag{40}$$

$$\begin{aligned}
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| s(1-t) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| t(1-s) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| (1-s)(1-t) ds dt, \\
|I_{14}| & \leq \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Lambda_1(t)| |\Delta_2(s)| ts ds dt
\end{aligned} \tag{41}$$

$$\begin{aligned}
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Lambda_1(t)| |\Delta_2(s)| s(1-t) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Lambda_1(t)| |\Delta_2(s)| t(1-s) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 |\Lambda_1(t)| |\Delta_2(s)| (1-s)(1-t) ds dt,
\end{aligned} \tag{42}$$

$$\begin{aligned}
|I_{15}| & \leq \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Lambda_2(s)| ts ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Lambda_2(s)| s(1-t) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Lambda_2(s)| t(1-s) ds dt \\
& + \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} |\Delta_1(t)| |\Lambda_2(s)| (1-s)(1-t) ds dt,
\end{aligned} \tag{42}$$

and

$$|I_{16}| \leq \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| tsdsdt \quad (43)$$

$$\begin{aligned} &+ \left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| s(1-t) dsdt \\ &+ \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| t(1-s) dsdt \\ &+ \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 |\Delta_1(t)| |\Delta_2(s)| (1-s)(1-t) dsdt. \end{aligned}$$

With the help of the inequalities (28)-(43), we get

$$\begin{aligned} &|I_1| + |I_2| + \cdots + |I_{16}| \quad (44) \\ &\leq [A_1 + A_2] [B_1 + B_2] \left(\left| \frac{\partial^2 f}{\partial t \partial s} (a, c) \right| + \left| \frac{\partial^2 f}{\partial t \partial s} (b, c) \right| + \left| \frac{\partial^2 f}{\partial t \partial s} (a, d) \right| + \left| \frac{\partial^2 f}{\partial t \partial s} (b, d) \right| \right). \end{aligned}$$

If the inequality (44) is written into (27), then we get desired result. This finish the proof of Theorem 3.2. \square

Remark 3.3. In Theorem 3.2, if we assign $\varphi(t) = t$ and $\psi(s) = s$ for all $(t, s) \in \Omega$, then Theorem 3.2 reduces to [23, Theorem 2].

Remark 3.4. In Theorem 3.2, let us now note that $\varphi(t) = \frac{t^\alpha}{\Gamma(\alpha)}$ and $\psi(s) = \frac{s^\beta}{\Gamma(\beta)}$ for all $(t, s) \in \Omega$ and $\alpha, \beta > 0$. Then, Theorem 3.2 reduces to [43, Theorem 2.1].

Corollary 3.5. In Theorem 3.2, if we choose $\varphi(t) = \frac{t^\alpha}{k\Gamma_k(\alpha)}$ and $\psi(s) = \frac{s^\beta}{k\Gamma_k(\beta)}$ for all $(t, s) \in \Omega$ and $\alpha, \beta, k > 0$, then we obtain the following inequality

$$\left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \left[\frac{\Gamma_k(\alpha+k)}{2(b-a)^{\frac{\alpha}{k}}} \left(J_{a+,k}^\alpha f\left(b, \frac{c+d}{2}\right) + J_{b-,k}^\alpha f\left(a, \frac{c+d}{2}\right) \right) \right] \right.$$

$$\left. + \frac{\Gamma_k(\beta+k)}{2(d-c)^{\frac{\beta}{k}}} \left[J_{c+,k}^\beta f\left(\frac{a+b}{2}, d\right) + J_{d-,k}^\beta f\left(\frac{a+b}{2}, c\right) \right] \right]$$

$$+ \frac{\Gamma_k(\alpha+k)\Gamma_k(\beta+k)}{4(b-a)^{\frac{\alpha}{k}}(d-c)^{\frac{\beta}{k}}}$$

$$\times \left[J_{a+,c+}^{\alpha,\beta,k} f(b, d) + J_{a+,d-}^{\alpha,\beta,k} f(b, c) + J_{b-,c+}^{\alpha,\beta,k} f(a, d) + J_{b-,d-}^{\alpha,\beta,k} f(a, c) \right]$$

$$\leq \frac{(b-a)(d-c)}{4} \left(\frac{1}{2} + \frac{k(1-2^{\frac{\alpha}{k}})}{2^{\frac{\alpha}{k}}(\alpha+k)} \right) \left(\frac{1}{2} + \frac{k(1-2^{\frac{\beta}{k}})}{2^{\frac{\beta}{k}}(\beta+k)} \right)$$

$$\times \left(\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right| + \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right| + \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right| + \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right| \right).$$

Theorem 3.6. Assume $f : \Omega \rightarrow \mathbb{R}$ is a partial differentiable mapping on Ω . Assume also $\left| \frac{\partial^2 f}{\partial t \partial s} \right|^q$, $q > 1$ is a convex function on the co-ordinates on Ω . Then, one obtains the following inequality

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \frac{1}{2\Lambda_1(1)} \left[{}_{a+}I_{\varphi}f\left(b, \frac{c+d}{2}\right) + {}_{b-}I_{\varphi}f\left(a, \frac{c+d}{2}\right) \right] \right. \\ & \quad \left. - \frac{1}{2\Lambda_2(1)} \left[{}_{c+}I_{\psi}f\left(\frac{a+b}{2}, d\right) + {}_{d-}I_{\psi}f\left(\frac{a+b}{2}, c\right) \right] \right. \\ & \quad \left. + \frac{1}{4\Lambda_1(1)\Lambda_2(1)} \left[{}_{a+, c+}I_{\varphi, \psi}f(b, d) + {}_{a+, d-}I_{\varphi, \psi}f(b, c) + {}_{b-, c+}I_{\varphi, \psi}f(a, d) + {}_{b-, d-}I_{\varphi, \psi}f(a, c) \right] \right|^{\frac{1}{q}} \\ & \leq \frac{(b-a)(d-c)}{4\Lambda_1(1)\Lambda_2(1)} [C_1 + C_2][D_1 + D_2] \\ & \times \left\{ \left(\frac{\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 3\left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 3\left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 9\left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\frac{3\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 9\left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 3\left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\frac{3\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 9\left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 3\left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\frac{9\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 3\left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 3\left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}} \right\}, \end{aligned}$$

where

$$\begin{cases} C_1 = \left(\int_0^{\frac{1}{2}} |\Lambda_1(t)|^p dt \right)^{\frac{1}{p}}, & D_1 = \left(\int_0^{\frac{1}{2}} |\Lambda_2(s)|^p ds \right)^{\frac{1}{p}}, \\ C_2 = \left(\int_{\frac{1}{2}}^1 |\Delta_1(t)|^p dt \right)^{\frac{1}{p}}, & D_2 = \left(\int_{\frac{1}{2}}^1 |\Delta_2(s)|^p ds \right)^{\frac{1}{p}}, \end{cases}$$

and $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. With the help of the Hölder's inequality for double integrals and by using the co-ordinated convexity

of $\left| \frac{\partial^2 f}{\partial t \partial s} \right|^q$ function on Ω , we have

$$|I_1| \leq \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)| |\Lambda_2(s)| \left| \frac{\partial^2 f}{\partial t \partial s} (ta + (1-t)b, sc + (1-s)d) \right| ds dt \quad (45)$$

$$\begin{aligned} & \leq \left(\int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} |\Lambda_1(t)|^p |\Lambda_2(s)|^p ds dt \right)^{\frac{1}{p}} \\ & \quad \times \left(\int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} \left| \frac{\partial^2 f}{\partial t \partial s} (ta + (1-t)b, sc + (1-s)d) \right|^q ds dt \right)^{\frac{1}{q}} \\ & \leq C_1 \cdot D_1 \left(\int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} \left\{ ts \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + s(1-t) \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q \right. \right. \\ & \quad \left. \left. + t(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + (1-t)(1-s) \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q \right\} ds dt \right)^{\frac{1}{q}} \\ & \leq C_1 \cdot D_1 \left(\frac{\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \end{aligned}$$

$$|I_2| \leq C_1 \cdot D_2 \left(\frac{3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (46)$$

$$|I_3| \leq C_2 \cdot D_1 \left(\frac{3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (47)$$

$$|I_4| \leq C_2 \cdot D_2 \left(\frac{9 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (48)$$

$$|I_5| \leq C_1 \cdot D_1 \left(\frac{9 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (49)$$

$$|I_6| \leq C_1 \cdot D_2 \left(\frac{3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (50)$$

$$|I_7| \leq C_2 \cdot D_1 \left(\frac{3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (51)$$

$$|I_8| \leq C_2 \cdot D_2 \left(\frac{\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (52)$$

$$|I_9| \leq C_1 \cdot D_1 \left(\frac{3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (53)$$

$$|I_{10}| \leq C_1 \cdot D_2 \left(\frac{\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (54)$$

$$|I_{11}| \leq C_2 \cdot D_1 \left(\frac{9 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (55)$$

$$|I_{12}| \leq C_2 \cdot D_2 \left(\frac{3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (56)$$

$$|I_{13}| \leq C_1 \cdot D_1 \left(\frac{3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (57)$$

$$|I_{14}| \leq C_1 \cdot D_2 \left(\frac{9 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (58)$$

$$|I_{15}| \leq C_2 \cdot D_1 \left(\frac{\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}, \quad (59)$$

and

$$|I_{16}| \leq C_2 \cdot D_2 \left(\frac{3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}}. \quad (60)$$

If the inequalities (45)-(60) are written into (27), then we obtain desired result. This is the end of the proof of Theorem 3.6. \square

Remark 3.7. In Theorem 3.6, let us consider that $\varphi(t) = \frac{t^\alpha}{\Gamma(\alpha)}$ and $\psi(s) = \frac{s^\beta}{\Gamma(\beta)}$ for all $(t, s) \in \Omega$. Then, Theorem 3.6 reduces to [43, Theorem 2.2].

Corollary 3.8. In Theorem 3.6, if we take $\varphi(t) = \frac{t^\frac{\alpha}{k}}{k\Gamma_k(\alpha)}$ and $\psi(s) = \frac{s^\frac{\beta}{k}}{k\Gamma_k(\beta)}$ for all $(t, s) \in \Omega$, then we get the following inequality

$$\begin{aligned} & \left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) - \left[\frac{\Gamma_k(\alpha+k)}{2(b-a)^\frac{\alpha}{k}} \left[J_{a+,k}^\alpha f\left(b, \frac{c+d}{2}\right) + J_{b-,k}^\alpha f\left(a, \frac{c+d}{2}\right) \right] \right. \right. \\ & \quad \left. \left. + \frac{\Gamma_k(\beta+k)}{2(d-c)^\frac{\beta}{k}} \left[J_{c+,k}^\beta f\left(\frac{a+b}{2}, d\right) + J_{d-,k}^\beta f\left(\frac{a+b}{2}, c\right) \right] \right] \right| \\ & \quad + \frac{\Gamma_k(\alpha+k)\Gamma_k(\beta+k)}{4(b-a)^\frac{\alpha}{k}(d-c)^\frac{\beta}{k}} \\ & \quad \times \left[J_{a+,c+}^{\alpha,\beta,k} f(b, d) + J_{a+,d-}^{\alpha,\beta,k} f(b, c) + J_{b-,c+}^{\alpha,\beta,k} f(a, d) + J_{b-,d-}^{\alpha,\beta,k} f(a, c) \right] \\ & \leq \frac{(b-a)(d-c)}{4} \left[\left(\frac{1}{2} + \frac{1-2^{p\frac{\alpha}{k}+1}}{2^{p\frac{\alpha}{k}+1}(p\frac{\alpha}{k}+1)} \right)^{\frac{1}{p}} + \left(\frac{1}{2^{p\frac{\alpha}{k}+1}(p\frac{\alpha}{k}+1)} \right)^{\frac{1}{p}} \right] \\ & \quad \times \left[\left(\frac{1}{2} + \frac{1-2^{p\frac{\beta}{k}+1}}{2^{p\frac{\beta}{k}+1}(p\frac{\beta}{k}+1)} \right)^{\frac{1}{p}} + \left(\frac{1}{2^{p\frac{\beta}{k}+1}(p\frac{\beta}{k}+1)} \right)^{\frac{1}{p}} \right] \\ & \quad \times \left\{ \left(\frac{9 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}} \right. \\ & \quad + \left. \left(\frac{\left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}} \right. \\ & \quad + \left. \left(\frac{3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}} \right. \\ & \quad + \left. \left(\frac{3 \left| \frac{\partial^2 f}{\partial t \partial s}(a, c) \right|^q + 9 \left| \frac{\partial^2 f}{\partial t \partial s}(b, c) \right|^q + \left| \frac{\partial^2 f}{\partial t \partial s}(a, d) \right|^q + 3 \left| \frac{\partial^2 f}{\partial t \partial s}(b, d) \right|^q}{64} \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

4. Conclusions

In this manuscript, we consider generalized fractional integrals to get midpoint type inequalities for the co-ordinated convex functions. Firstly, it is established an identity for twice partially differentiable mappings. By using the this identity, some midpoint type inequalities via generalized fractional integrals are proved. Furthermore, the main results reduce some midpoint inequalities given in earlier works for Riemann integrals and Riemann-Liouville fractional integrals. In addition to this, it is introduced some new inequalities for k -Riemann-Liouville fractional integrals as special cases of our results. We propose for forthcoming researchers that the methods and techniques used in this study can be established similar inequalities for different kinds of co-ordinated convexity.

References

- [1] B. Ahmad, A. Alsaedi, M. Kirane, and B. T. Torebek, *Hermite-Hadamard, Hermite-Hadamard-Fejer, Dragomir-Agarwal and Pachpatte type inequalities for convex functions via new fractional integrals*, Journal of Computational and Applied Mathematics, vol. 353, pp. 12
- [2] G. Anastassiou, *General fractional hermite-hadamard inequalities using m -convexity and (s, m) -convexity*, Frontiers in Time Scales and Inequalities, vol. 237, p.
- [3] M. K. Bakula, M. E. Ozdemir and J. Pečarić, *Hadamard type inequalities for m -convex and (α, m) -convex functions*, J. Inequal. Pure Appl. Math 9.4, 2008, 12.
- [4] H. Budak, F. Hezenci, and H. Kara, *On parameterized inequalities of Ostrowski and Simpson type for convex functions via generalized fractional integrals*, Mathematical Methods in the Applied Sciences, 44(17), 2021, 12522-12536.
- [5] H. Budak, F. Hezenci, and H. Kara, *On generalized Ostrowski, Simpson and Trapezoidal type inequalities for co-ordinated convex functions via generalized fractional integrals*, Advances in Difference Equations, 2021, 2021, 1-32.
- [6] H. Budak, H. Kara and R. Kapucu, *New midpoint type inequalities for generalized fractional integral*, Computational Methods for Differential Equations, 2021.
- [7] H. Budak, E. Pehlivan and P. Kösem, *On new extensions of Hermite-Hadamard inequalities for generalized fractional integrals*. Sahand Communications in Mathematical Analysis, 2021.
- [8] H. Budak and P. Agarwal, *New generalized midpoint type inequalities for fractional integral*, Miskolc Math. Notes, 20(2019), 781-793.
- [9] F. Chen, *On Hermite-Hadamard type Inequalities for s -convex functions on the coordinates via Riemann-Liouville fractional integrals*, Journal of Applied Mathematics, Volume 2014 (2014), Article ID 248710, 8 pages.
- [10] F. Chen, *A note on the Hermite-Hadamard inequality for convex functions on the co-ordinates*, J. Math. Inequal., 8(4), (2014), 915-923.
- [11] M. R. Delavar, *New bounds for Hermite-Hadamard's trapezoid and mid-point type inequalities via fractional integrals*. Miskolc Mathematical Notes, 20(2), 2019, 849-861.
- [12] S.S. Dragomir and R. P. Agarwal, *Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula*, Appl. Math. lett. 11 (5) (1998) 91–95.
- [13] S. S. Dragomir, *On Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane*, Taiwanese J Math., 4 (2001), 775–788.
- [14] F. Ertuğral and M. Z. Sarikaya, *Simpson type integral inequalities for generalized fractional integral*, Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas, 113(4), 2019, 3115-3124.
- [15] J. Han, P. O. Mohammed and H. Zeng, *Generalized fractional integral inequalities of Hermite-Hadamard-type for a convex function*, Open Mathematics, 18(1), 2020, 794-806.
- [16] F. Hezenci, H. Budak, and H. Kara, *New version of fractional Simpson type inequalities for twice differentiable functions*, Advances in Difference Equations, 2021, 2021, 1-10.
- [17] M. Iqbal, M.I. Bhatti, K. Nazeer, *Generalization of inequalities analogous to Hermite-Hadamard inequality via fractional integrals*, Bull. Korean Math. Soc. 52 (2015), no. 3, 707-716.
- [18] A. Kashuri and R. Liko, *On Fejér type inequalities for convex mappings utilizing generalized fractional integrals*, Applications & Applied Mathematics, 2020, 15(1), 135-150.
- [19] H. Kavurmacı, M. Avci, and M. E. Ozdemir, *New inequalities of Hermite-Hadamard type for convex functions with applications*, Journal of Inequalities and Applications, 2011(1), 1-11.
- [20] U. S. Kirmaci, *Inequalities for differentiable mappings and applications to special means of real numbers to midpoint formula*, Appl. Math. Comput., vol. 147, no. 5, pp. 137–146, 2004.
- [21] U. S. Kirmaci, M. K. Bakula, M. K., M. E. Özdemir and J. Pečarić, *Hadamard-type inequalities for s -convex functions*, Applied Mathematics and Computation, 193(1), 2007, 26-35.
- [22] M. A. Latif and M. Alomari, *On the Hadamard-type inequalities for h -convex functions on the co-ordinates*, Int. J. of Math. Analysis, 3(33), 2009, 1645-1656.
- [23] M. A. Latif and S. S. Dragomir, *On some new inequalities for differentiable coordinated convex functions*, Journal of Inequalities and Applications, Vol. 2012, 2012, No. 28, pp. 1-13.
- [24] M. V. Marcela, *New inequalities for co-ordinated convex functions via Riemann-Liouville fractional calculus*, Tamkang Journal of Mathematics, 45(3), 2014, 285-296.
- [25] P. O. Mohammed, *On new trapezoid type inequalities for h -convex functions via generalized fractional integral*, Turkish J. Anal. Number Theory 2018, 6, 125–128.
- [26] P. O. Mohammed and M. Z. Sarikaya, *On generalized fractional integral inequalities for twice differentiable convex functions*, J. Comput. Appl. Math. 2020, 372, 112740.
- [27] P. O. Mohammed, *Hermite-Hadamard inequalities for Riemann-Liouville fractional integrals of a convex function with respect to a monotone function*, Math. Meth. Appl. Sci. 2019, 1-11.
- [28] M. A. Noor, K. I. Noor, M. U. Awan and J. Li, *On Hermite-Hadamard inequalities for h -Preinvex functions*, Filomat, 28(7), 2014, 1463-1474.
- [29] M. A. Noor and M. U. Awan, *Some integral inequalities for two kinds of convexities via fractional integrals*, Trans. J. Math. Mech, vol. 5, no. 2, pp. 129-136, 2013.
- [30] M.E. Özdemir, E. Set and M.Z. Sarikaya, *New some Hadamard's type inequalities for co-ordinated m -convex and (α, m) -convex functions*, Hacettepe Journal of Mathematics and Statistics, 40(2), 2011, 219-229.
- [31] M. E. Ozdemir, C. Yıldız and A. O. Akdemir, *On some new Hadamard-type inequalities for co-ordinated quasi-convex functions*, Hacettepe Journal of Mathematics and Statistics, 41(5), 2012, 697-707.
- [32] M. E. Ozdemir, M. Avci-Ardıç, and H. Kavurmacı- Onalan, *Hermite-hadamard type inequalities for s -convex and s -concave functions via fractional integrals*, Turkish Journal of Science, 1(1), pp. 28-40, 2016.

- [33] S. Qaisar and S. Hussain, *On hermite-hadamard type inequalities for functions whose first derivative absolute values are convex and concave*, *Fasciculi Mathematici*, 58(1), 2017, 155–166.
- [34] F. Qi, P. O. Mohammed J.-C. Yao and Y.-H. Yao, *Generalized fractional integral inequalities of Hermite–Hadamard type for (α, m) -convex functions*, *J Inequal Appl* 2019, 135 (2019).
- [35] M. Z. Sarikaya, A. Saglam, and H. Yildirim, *On some Hadamard-type inequalities for h-convex functions*, *J. Math. Inequal.*, 2(3), 2008, 335–34.
- [36] M. Z. Sarikaya and H. Yildirim, *On Hermite-Hadamard type inequalities for Riemann-Liouville fractional integrals*, *Miskolc Mathematical Notes*, 17(2), 2016, pp. 1049-1059.
- [37] M.Z. Sarikaya and H. Budak, *Generalized Hermite-Hadamard type integral inequalities for fractional integrals*, *Filomat*, 30(2016), 1315–1326
- [38] M. Z. Sarikaya, E. Set, M. E. Ozdemir and S. S. Dragomir, *New some Hadamard's type inequalities for coordinated convex functions*, *Tamsui Oxford Journal of Information and Mathematical Sciences*, 28(2), 2012, 137–152.
- [39] M. Z. Sarikaya, E. Set, H. Yaldiz, N. Basak, *Hermite–Hadamard's inequalities for fractional integrals and related fractional inequalities*, *Math Comput Model*, 57 (9–10) (2013), 2403–2407.
- [40] M. Z. Sarikaya, *On the Hermite-Hadamard-type inequalities for co-ordinated convex function via fractional integrals*, *Integral Transforms and Special Functions*, 25(2), 2014, 134–147.
- [41] M. Z. Sarikaya, F. Ertugral, *On the generalized Hermite-Hadamard inequalities*, *Annals of the University of Craiova, Mathematics and Computer Science Series*, 47 (1) (2020), 193–213.
- [42] M. Z. Sarikaya, A. Akkurt, H. Budak, M. E. Yildirim, H. Yildirim, *Hermite-Hadamard's inequalities for conformable fractional integrals*, *Konuralp Journal of Mathematics*, 8 (2) (2020) 376–383.
- [43] T. Tunç , M. Z. Sarikaya, H. Yaldiz, *Fractional Hermite-Hadamard's type inequality for the co-ordinated convex functions*, *TWMS J. Pure Appl. Math.*, Vol. 11, 2020, No. 1, pp. 3-29.
- [44] M. E. Turkyay, M.Z. Sarikaya , H. Budak, and H. Yildirim, *Some Hermite-Hadamard type inequalities for co-ordinated convex functions via generalized fractional integrals*, Submitted, ResearchGate Article: <https://www.researchgate.net/publication/321803898>.
- [45] B.-Y. Xi, R.-F. Bai and F. Qi, *Hermite–Hadamard type inequalities for the m-and (α, m) -geometrically convex functions*, *Aequationes mathematicae* 84.3, 2012, 261-269.
- [46] Y. Zhang and J. Wang, *On some new Hermite-Hadamard inequalities involving Riemann-Liouville fractional integrals*, *Journal of Inequalities and Applications*, vol. 2013, no. 1, 2013, p. 220.
- [47] D. Zhao, M. A. Ali,A. Kashuri, H. Budak M. Z. Sarikaya, *Hermite–Hadamard-type inequalities for the interval-valued approximately h-convex functions via generalized fractional integrals*, *Journal of Inequalities and Applications*, 2020(1), 1-38.