



Probabilistic Lah numbers and Lah-Bell polynomials

Siqi Dong^{a,*}, Yuankui Ma^a, Taekyun Kim^{a,b}, Dae San Kim^c, Wenpeng Zhang^{a,d}

^a*School of Science, Xi'an Technological University, Xi'an 710021, Shaanxi, China*

^b*Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea*

^c*Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea*

^d*School of mathematics, Northwest University, Xi'an 710127, Shaanxi, China*

Abstract. Let Y be a random variable whose moment generating function exists in some neighborhood of the origin. The aim of this paper is to study the probabilistic Lah numbers associated with Y and the probabilistic Lah-Bell polynomials associated with Y , as probabilistic versions of the Lah numbers and the Lah-Bell polynomials, respectively. We derive some properties, explicit expressions, recurrence relations and certain identities for those numbers and polynomials. In addition, we treat the special cases that Y is the Poisson random variable with parameter $\alpha > 0$ and the Bernoulli random variable with probability of success p .

1. Introduction

The Stirling number of the second kind $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ (see (7)) enumerates the number of ways a set of n elements can be partitioned into k nonempty subsets, while the n th Bell number $\phi_n = \phi_n(1)$ (see (6)) counts the number of ways a set of n elements can be partitioned into nonempty subsets. The classical unsigned Lah number $L(n, k)$ (see (3), (4), (5)) counts the number of ways a set of n elements can be partitioned into k nonempty linearly ordered subsets. Just as the Bell number is to the Stirling numbers, the Lah-Bell number is to the unsigned Lah numbers. More precisely, the n th Lah-Bell number $B_n^L = B_n^L(1)$ (see (8), (9)) is defined as the number of ways a set of n elements can be partitioned into nonempty linearly ordered subsets.

Recently, Kim et al. proposed applying probabilistic methods to various generalized polynomials. Kim and Kim (see [17]) defined probabilistic bivariate Bell polynomials associated with Y , and this framework was applied to other polynomials, including probabilistic degenerate derangement polynomials associated with Y (see [18]) and Dowling polynomials associated with Y (see [25]). Concurrently, Khan and Ahmad have been exploring a generalization that combines both probabilistic and degenerate parameters. This includes the study of probabilistic degenerate Jindalrae-Stirling polynomials (see [11]) and Bell-based partially degenerate Genocchi polynomials (see [5]), often with applications in number theory and analysis.

2020 *Mathematics Subject Classification.* Primary 11B73; Secondary 11B83.

Keywords. probabilistic Lah numbers; probabilistic Lah-Bell polynomials

Received: 04 January 2025; Accepted: 18 August 2025

Communicated by Paola Bonacini

Research supported by National Natural Science Foundation of China (No.12271320).

* Corresponding author: Siqi Dong

Email addresses: dongsiqi@st.xatu.edu.cn (Siqi Dong), mayuankui@xatu.edu.cn (Yuankui Ma), kimtk2015@gmail.com (Taekyun Kim), dskim@sogang.ac.kr (Dae San Kim), wpzhang888@163.com (Wenpeng Zhang)

ORCID iDs: <https://orcid.org/0009-0004-9863-3128> (Siqi Dong), <https://orcid.org/0000-0002-5556-0633> (Yuankui Ma), <https://orcid.org/0000-0002-3731-7023> (Taekyun Kim), <https://orcid.org/0000-0001-9599-7015> (Dae San Kim), <https://orcid.org/0000-0002-2560-3559> (Wenpeng Zhang)

Based on the framework established by Kim et al., we defined probabilistic Lah numbers and probabilistic Lah Bell polynomials and studied their properties.

The aim of this paper is to study probabilistic versions of the Lah numbers $L(n, k)$, and the Lah-Bell polynomials $B_n^L(x)$, which are a natural extension of Lah-Bell numbers, namely the probabilistic Lah numbers associated with Y and the probabilistic Lah-Bell polynomials associated with Y . Here Y is a random variable whose moment generating function exists in some neighborhood of the origin. We derive some properties, explicit expressions, recurrence relations and certain identities for those numbers and polynomials. In addition, we treat the special cases that Y is the Poisson random variable with parameter $\alpha > 0$ and the Bernoulli random variable with probability of success p .

The outline of this paper is as follows. In Section 1, we recall the Stirling numbers of the second kind and the Bell polynomials. We remind the reader of the Lah numbers and the Lah-Bell polynomials. We recall the partial Bell polynomials and the complete Bell polynomials. Section 2 is the main result of this paper. Assume that Y is a random variable such that the moment generating function of Y , $E[e^{tY}] = \sum_{n=0}^{\infty} \frac{t^n}{n!} E[Y^n]$, ($|t| < r$), exists for some $r > 0$. Let $(Y_j)_{j \geq 1}$ be a sequence of mutually independent copies of the random variable Y , and let $S_k = Y_1 + Y_2 + \cdots + Y_k$, ($k \geq 1$), with $S_0 = 0$. Then we first define the probabilistic Lah numbers associated with the random variable Y , $L_Y(n, k)$, which are probabilistic versions of the Lah numbers and reduce to those numbers for $Y = 1$. In Theorem 2.1, we derive an expression of $L_Y(n, k)$ in terms of $E[\langle S_l \rangle_n]$, ($l = 0, \dots, k$). In Theorem 2.2, we show that $L_Y(n, k) = \frac{1}{k!} E[\Delta_{Y_1, Y_2, \dots, Y_k} \langle 0 \rangle_n]$, (see (17)). Next, we define the probabilistic Lah-Bell polynomials associated with the random variable Y , $B_n^{(L, Y)}(x)$, which are probabilistic versions of Lah-Bell polynomials. We obtain in Theorem 2.3 the generating function of the polynomials $B_n^{(L, Y)}(x)$, from which a Dobinski-like formula for those ones are deduced in Theorem 2.4. Two expressions of $B_n^{(L, Y)}(x)$ are obtained in terms of the partial Bell polynomials in Theorems 2.5 and 2.8. A recurrence relation for $B_n^{(L, Y)}(x)$ is derived in Theorem 2.6. It is shown that $B_n^{(L, Y)}(x)$ satisfies the binomial identity in Theorem 2.7. Two identities involving $L_Y(n, k)$ and $B_n^{(L, Y)}(x)$ are obtained in Theorems 2.9 and 2.10. A finite sum expression for the k th derivative of $B_n^{(L, Y)}(x)$ is shown in Theorem 2.11. Let Y be the Poisson random variable with parameter $\alpha > 0$. Then we show $E[\langle S_k \rangle_n] = B_n^L(\alpha k)$ in Theorem 2.12 and in terms of the Lah numbers and the Bell polynomials a finite sum expression of $B_n^{(L, Y)}(x)$ in Theorem 2.13. Finally, in Theorem 2.14 we show $B_n^{(L, Y)}(x) = B_n^L(px)$ and $L_Y(n, k) = p^k L(n, k)$ when Y is the Bernoulli random variable with probability of success p .

We recall that the falling factorial sequence is defined by

$$(x)_0 = 1, \quad (x)_n = x(x-1)(x-2) \cdots (x-n+1), \quad (n \geq 1), \quad (\text{see}[1-30]). \quad (1)$$

Also, the rising factorial sequence is given by

$$\langle x \rangle_0 = 1, \quad \langle x \rangle_n = x(x+1)(x+2) \cdots (x+n-1), \quad (n \geq 1). \quad (2)$$

For any integers n, k with $n \geq k \geq 0$, the unsigned Lah numbers are defined by

$$\langle x \rangle_n = \sum_{k=0}^n L(n, k) (x)_k, \quad (n \geq 0), \quad (\text{see}[10, 14, 15, 26]). \quad (3)$$

Thus, by (3), we get

$$\frac{1}{k!} \left(\frac{1}{1-t} - 1 \right)^k = \sum_{n=k}^{\infty} L(n, k) \frac{t^n}{n!}, \quad (k \geq 0), \quad (\text{see}[13-15, 19, 20, 22, 23]). \quad (4)$$

From (4), we note that

$$L(n, k) = \frac{n!}{k!} \binom{n-1}{k-1}, \quad (n \geq k \geq 0), \quad (\text{see}[10, 14, 15, 27]). \quad (5)$$

It is well known that the Bell polynomials are defined by

$$\phi_n(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k, \quad (\text{see}[16, 27]). \quad (6)$$

where $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}$ are the Stirling numbers of the second kind given by

$$x^n = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} (x)_k, \quad (\text{see}[16, 21, 24, 26, 27]). \quad (7)$$

In view of (6), Lah-Bell polynomials are define by

$$B_n^L(x) = \sum_{l=0}^n L(n, l) x^l, \quad (n \geq 0), \quad (\text{see}[14, 15]). \quad (8)$$

Thus, by (4) and (8), we get

$$e^{x(\frac{1}{1-t}-1)} = \sum_{n=0}^{\infty} B_n^L(x) \frac{t^n}{n!}, \quad (\text{see}[13 - 16, 19 - 24]). \quad (9)$$

For $k \geq 0$, the partial Bell polynomials are defined by

$$\frac{1}{k!} \left(\sum_{j=1}^{\infty} x_j \frac{t^j}{j!} \right)^k = \sum_{n=k}^{\infty} B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) \frac{t^n}{n!}, \quad (\text{see}[26]). \quad (10)$$

Thus, by (10), we get

$$B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{k_1+k_2+\dots+k_{n-k+1}=k \\ k_1+2k_2+\dots+(n-k+1)k_{n-k+1}=n}} \frac{n!}{k_1!k_2!\dots k_{n-k+1}!} \left(\frac{x_1}{1!} \right)^{k_1} \left(\frac{x_2}{2!} \right)^{k_2} \dots \left(\frac{x_{n-k+1}}{(n-k+1)!} \right)^{k_{n-k+1}}. \quad (11)$$

The complete Bell polynomials are given by

$$\exp \left(\sum_{j=1}^{\infty} x_j \frac{t^j}{j!} \right) = \sum_{n=0}^{\infty} B_n(x_1, x_2, \dots, x_n) \frac{t^n}{n!}, \quad (\text{see}[10, 20, 23]). \quad (12)$$

Thus, by (10) and (11), we get

$$B_n(x_1, x_2, \dots, x_n) = \sum_{k=0}^n B_{n,k}(x_1, x_2, \dots, x_{n-k+1}), \quad (\text{see}[10, 15, 20]). \quad (13)$$

2. Probabilistic Lah numbers and Lah-Bell polynomials

Throughout this section, we assume that Y is a random variable such that the moment generating function of Y , $E[e^{tY}] = \sum_{n=0}^{\infty} E[Y^n] \frac{t^n}{n!}$, ($|t| < r$), exists for some $r > 0$, (see [3,16]). We let $(Y_j)_{j \geq 1}$ be a sequence of mutually independent copies of the random variable Y , and let

$$S_0 = 0, \quad S_k = Y_1 + Y_2 + \dots + Y_k, \quad (k \in \mathbb{N}), \quad (\text{see}[3, 20]). \quad (14)$$

We define the probabilistic Lah numbers associated with the random variable Y by

$$\frac{1}{k!} \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)^k = \sum_{n=k}^{\infty} L_Y(n, k) \frac{t^n}{n!}, \quad (k \geq 0). \quad (15)$$

When $Y = 1$, we note that $L_Y(n, k) = L(n, k)$. From (15), we note that

$$\begin{aligned} \sum_{n=k}^{\infty} L_Y(n, k) \frac{t^n}{n!} &= \frac{1}{k!} \sum_{l=0}^k \binom{k}{l} (-1)^{k-l} E \left[\left(\frac{1}{1-t} \right)^{Y_1+Y_2+\dots+Y_l} \right] \\ &= \frac{1}{k!} \sum_{l=0}^k \binom{k}{l} (-1)^{k-l} E \left[\left(\frac{1}{1-t} \right)^{S_l} \right] = \sum_{n=0}^{\infty} \frac{1}{k!} \sum_{l=0}^k \binom{k}{l} (-1)^{k-l} E[\langle S_l \rangle_n] \frac{t^n}{n!}. \end{aligned} \quad (16)$$

Therefore, by comparing the coefficients on both sides of (16), we obtain the following theorem.

Theorem 2.1. For any integers n, k with $n \geq k \geq 0$, we have

$$L_Y(n, k) = \frac{1}{k!} \sum_{l=0}^k \binom{k}{l} (-1)^{k-l} E[\langle S_l \rangle_n].$$

The operators Δ_y and $\Delta_{y_1, y_2, \dots, y_m}$ are respectively given by

$$\Delta_y f(x) = f(x+y) - f(x), \quad \Delta_{y_1, y_2, \dots, y_m} f(x) = \Delta_{y_1} \circ \Delta_{y_2} \circ \Delta_{y_3} \circ \dots \circ \Delta_{y_m} f(x), \quad (17)$$

where $y \in \mathbb{R}$, and $(y_1, y_2, \dots, y_m) \in \mathbb{R}^m$. Then, by (17), we get

$$\begin{aligned} \Delta_{y_1, y_2, \dots, y_m} \left(\frac{1}{1-t} \right)^x &= \left(\left(\frac{1}{1-t} \right)^{y_1} - 1 \right) \left(\left(\frac{1}{1-t} \right)^{y_2} - 1 \right) \dots \left(\left(\frac{1}{1-t} \right)^{y_m} - 1 \right) \left(\frac{1}{1-t} \right)^x \\ &= \sum_{n=m}^{\infty} \frac{t^n}{n!} \Delta_{y_1, y_2, \dots, y_m} \langle x \rangle_n. \end{aligned} \quad (18)$$

From (15), (18), we note that

$$\begin{aligned} \frac{1}{m!} \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)^m &= \frac{1}{m!} \left(E \left[\left(\frac{1}{1-t} \right)^Y - 1 \right] \right)^m \\ &= \frac{1}{m!} \left(E \left[\left(\left(\frac{1}{1-t} \right)^{Y_1} - 1 \right) \left(\left(\frac{1}{1-t} \right)^{Y_2} - 1 \right) \dots \left(\left(\frac{1}{1-t} \right)^{Y_m} - 1 \right) \right] \right) \\ &= \sum_{n=m}^{\infty} \frac{1}{m!} E[\Delta_{Y_1, Y_2, \dots, Y_m} \langle 0 \rangle_n] \frac{t^n}{n!}. \end{aligned} \quad (19)$$

Therefore, by (19), we obtain the following theorem.

Theorem 2.2. For any integers n, m with $n \geq m \geq 0$, we have

$$L_Y(n, m) = \frac{1}{m!} E[\Delta_{Y_1, Y_2, \dots, Y_m} \langle 0 \rangle_n].$$

We define the probabilistic Lah-Bell polynomials associated with the random variable Y by

$$B_n^{(L, Y)}(x) = \sum_{k=0}^n L_Y(n, k) x^k, \quad (n \geq 0). \quad (20)$$

When $Y = 1$, we have $B_n^{(L,Y)}(x) = B_n^L(x)$, (see (8)).

Form (20), we derive the generating function of the probabilistic Lah-Bell polynomials associated the random variable Y , which is given by

$$\begin{aligned} \sum_{n=0}^{\infty} B_n^{(L,Y)}(x) \frac{t^n}{n!} &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n L_Y(n, k) x^k \right) \frac{t^n}{n!}. \\ &= \sum_{k=0}^{\infty} x^k \left(\sum_{n=k}^{\infty} L_Y(n, k) \frac{t^n}{n!} \right) = \sum_{k=0}^{\infty} x^k \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)^k \frac{1}{k!} \\ &= \exp \left(x \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right) \right). \end{aligned} \quad (21)$$

Thus, by (21), we obtain the following theorem.

Theorem 2.3. *The generating function of the probabilistic Lah-Bell polynomials of the random variable Y is given by*

$$e^{x \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)} = \sum_{n=0}^{\infty} B_n^{(L,Y)}(x) \frac{t^n}{n!}. \quad (22)$$

By (22), we get

$$\begin{aligned} \sum_{n=0}^{\infty} B_n^{(L,Y)}(x) \frac{t^n}{n!} &= e^{x \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)} = e^{-x} e^{xE \left[\left(\frac{1}{1-t} \right)^Y \right]} \\ &= e^{-x} \sum_{k=0}^{\infty} \frac{x^k}{k!} \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] \right)^k = e^{-x} \sum_{k=0}^{\infty} \frac{x^k}{k!} E \left[\left(\frac{1}{1-t} \right)^{Y_1} \cdots \left(\frac{1}{1-t} \right)^{Y_k} \right] \\ &= e^{-x} \sum_{k=0}^{\infty} \frac{x^k}{k!} E \left[\left(\frac{1}{1-t} \right)^{S_k} \right] = \sum_{n=0}^{\infty} \left(e^{-x} \sum_{k=0}^{\infty} \frac{x^k}{k!} E[\langle S_k \rangle_n] \right) \frac{t^n}{n!}. \end{aligned} \quad (23)$$

Therefore, by (23), we obtain the following Dobinski-like formula.

Theorem 2.4. *For any integer $n \geq 0$, we have*

$$B_n^{(L,Y)}(x) = e^{-x} \sum_{k=0}^{\infty} \frac{x^k}{k!} E[\langle S_k \rangle_n].$$

From (22), we note that

$$\begin{aligned} \sum_{n=0}^{\infty} B_n^{(L,Y)}(x) \frac{t^n}{n!} &= e^{x \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)} \\ &= e^{x \sum_{j=1}^{\infty} E[\langle Y \rangle_j] \frac{t^j}{j!}} = \sum_{k=0}^{\infty} \frac{1}{k!} \left(x \sum_{j=1}^{\infty} E[\langle Y \rangle_j] \frac{t^j}{j!} \right)^k \\ &= \sum_{k=0}^{\infty} \sum_{n=k}^{\infty} B_{n,k} \left(x E[\langle Y \rangle_1], x E[\langle Y \rangle_2], \dots, x E[\langle Y \rangle_{n-k+1}] \right) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n B_{n,k} \left(x E[\langle Y \rangle_1], x E[\langle Y \rangle_2], \dots, x E[\langle Y \rangle_{n-k+1}] \right) \right) \frac{t^n}{n!}. \end{aligned} \quad (24)$$

Therefore, by comparing the coefficients on both sides of (24), we obtain the following theorem.

Theorem 2.5. For any integers n, k with $n \geq k \geq 0$, we have

$$B_n^{(L,Y)}(x) = \sum_{k=0}^n B_{n,k} \left(xE[\langle Y \rangle_1], xE[\langle Y \rangle_2], \dots, xE[\langle Y \rangle_{n-k+1}] \right).$$

Now, we observe that

$$\begin{aligned} \sum_{n=0}^{\infty} B_{n+1}^{(L,Y)}(x) \frac{t^n}{n!} &= \frac{d}{dt} \sum_{n=0}^{\infty} B_n^{(L,Y)}(x) \frac{t^n}{n!} = \frac{d}{dt} e^{x \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)} \\ &= x E \left[Y \left(\frac{1}{1-t} \right)^{Y+1} \right] e^{x \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)} = x \sum_{k=0}^{\infty} E[\langle Y \rangle_{k+1}] \frac{t^k}{k!} \sum_{m=0}^{\infty} B_m^{(L,Y)}(x) \frac{t^m}{m!} \\ &= \sum_{n=0}^{\infty} \left(x \sum_{k=0}^n \binom{n}{k} E[\langle Y \rangle_{k+1}] B_{n-k}^{(L,Y)}(x) \right) \frac{t^n}{n!}. \end{aligned} \quad (25)$$

Therefore, by (25), we obtain the following theorem.

Theorem 2.6. For any integer $n \geq 0$, we have the recurrence relation

$$B_{n+1}^{(L,Y)}(x) = x \sum_{k=0}^n \binom{n}{k} E[\langle Y \rangle_{k+1}] B_{n-k}^{(L,Y)}(x).$$

From (22), we have

$$\begin{aligned} \sum_{n=0}^{\infty} B_n^{(L,Y)}(x+y) \frac{t^n}{n!} &= e^{(x+y) \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)} = e^{x \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)} e^{y \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)} \\ &= \sum_{k=0}^{\infty} B_k^{(L,Y)}(x) \frac{t^k}{k!} \sum_{m=0}^{\infty} B_m^{(L,Y)}(y) \frac{t^m}{m!} = \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} B_k^{(L,Y)}(x) B_{n-k}^{(L,Y)}(y) \right) \frac{t^n}{n!}. \end{aligned} \quad (26)$$

Therefore, by (26), we obtain the following binomial identity.

Theorem 2.7. For any integer $n \geq 0$, we have

$$B_n^{(L,Y)}(x+y) = \sum_{k=0}^n \binom{n}{k} B_k^{(L,Y)}(x) B_{n-k}^{(L,Y)}(y).$$

By (22), we see that

$$\begin{aligned} \sum_{n=0}^{\infty} B_n^{(L,Y)}(x) \frac{t^n}{n!} &= e^{x \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)} \\ &= \sum_{k=0}^{\infty} \binom{x}{k} \left(e^{E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1} - 1 \right)^k = \sum_{k=0}^{\infty} k! \binom{x}{k} \frac{1}{k!} \left(\sum_{j=1}^{\infty} B_j^{(L,Y)}(1) \frac{t^j}{j!} \right)^k \\ &= \sum_{k=0}^{\infty} k! \binom{x}{k} \sum_{n=k}^{\infty} B_{n,k} \left(B_1^{(L,Y)}(1), B_2^{(L,Y)}(1), \dots, B_{n-k+1}^{(L,Y)}(1) \right) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n k! \binom{x}{k} B_{n,k} \left(B_1^{(L,Y)}(1), B_2^{(L,Y)}(1), \dots, B_{n-k+1}^{(L,Y)}(1) \right) \right) \frac{t^n}{n!}. \end{aligned} \quad (27)$$

Therefore, by (27), we obtain the following theorem.

Theorem 2.8. For any integer $n \geq 0$, we have

$$B_n^{(L,Y)}(x) = \sum_{k=0}^n k! \binom{x}{k} B_{n,k}^{(L,Y)}(1, B_2^{(L,Y)}(1), \dots, B_{n-k+1}^{(L,Y)}(1)).$$

Note that

$$te^{x\left(E\left[\left(\frac{1}{1-t}\right)^Y\right]-1\right)} = t \sum_{j=0}^{\infty} B_j^{(L,Y)}(x) \frac{t^j}{j!} = \sum_{j=1}^{\infty} j B_{j-1}^{(L,Y)}(x) \frac{t^j}{j!}. \quad (28)$$

By (28) and for $k \geq 0$, we get

$$\begin{aligned} \left(\sum_{j=1}^{\infty} j B_{j-1}^{(L,Y)}(x) \frac{t^j}{j!} \right)^k &= t^k e^{kx\left(E\left[\left(\frac{1}{1-t}\right)^Y\right]-1\right)} \\ &= t^k \sum_{j=0}^{\infty} k^j x^j \frac{1}{j!} \left(E\left[\left(\frac{1}{1-t}\right)^Y\right] - 1 \right)^j = t^k \sum_{j=0}^{\infty} k^j x^j \sum_{n=j}^{\infty} L_Y(n, j) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \sum_{j=0}^n k^j x^j L_Y(n, j) \frac{t^{n+k}}{n!} = \sum_{n=k}^{\infty} \sum_{j=0}^{n-k} k^j x^j L_Y(n-k, j) \frac{t^n}{(n-k)!} \\ &= \sum_{n=k}^{\infty} k! \sum_{j=0}^{n-k} \binom{n}{k} k^j x^j L_Y(n-k, j) \frac{t^n}{n!}. \end{aligned} \quad (29)$$

From (29), we have

$$\begin{aligned} \sum_{n=k}^{\infty} \sum_{j=0}^{n-k} \binom{n}{k} k^j x^j L_Y(n-k, j) \frac{t^n}{n!} &= \frac{1}{k!} \left(\sum_{j=1}^{\infty} j B_{j-1}^{(L,Y)}(x) \frac{t^j}{j!} \right)^k \\ &= \sum_{n=k}^{\infty} B_{n,k}^{(L,Y)}(B_0^{(L,Y)}(x), 2B_1^{(L,Y)}(x), \dots, (n-k+1)B_{n-k}^{(L,Y)}(x)) \frac{t^n}{n!}. \end{aligned} \quad (30)$$

Therefore, by (30), we obtain the following theorem.

Theorem 2.9. For any integers n, k with $n \geq k \geq 0$, we have

$$\begin{aligned} \sum_{j=0}^{n-k} \binom{n}{k} k^j x^j L_Y(n-k, j) \\ = B_{n,k}^{(L,Y)}(B_0^{(L,Y)}(x), 2B_1^{(L,Y)}(x), \dots, (n-k+1)B_{n-k}^{(L,Y)}(x)). \end{aligned}$$

Now, we observe that

$$\begin{aligned} \sum_{n=k}^{\infty} B_{n,k}^{(L,Y)}(B_1^{(L,Y)}(x), B_2^{(L,Y)}(x), \dots, B_{n-k+1}^{(L,Y)}(x)) \frac{t^n}{n!} \\ = \frac{1}{k!} \left(\sum_{j=1}^{\infty} B_j^{(L,Y)}(x) \frac{t^j}{j!} \right)^k = \frac{1}{k!} \left(e^{x\left(E\left[\left(\frac{1}{1-t}\right)^Y\right]-1\right)} - 1 \right)^k \\ = \sum_{j=k}^{\infty} \left\{ \begin{matrix} j \\ k \end{matrix} \right\} x^j \frac{1}{j!} \left(E\left[\left(\frac{1}{1-t}\right)^Y\right] - 1 \right)^j \\ = \sum_{j=k}^{\infty} \left\{ \begin{matrix} j \\ k \end{matrix} \right\} x^j \sum_{n=j}^{\infty} L_Y(n, j) \frac{t^n}{n!} = \sum_{n=k}^{\infty} \left(\sum_{j=k}^n \left\{ \begin{matrix} j \\ k \end{matrix} \right\} L_Y(n, j) x^j \right) \frac{t^n}{n!}. \end{aligned} \quad (31)$$

Therefore, by comparing the coefficients on both sides of (31), we obtain following theorem.

Theorem 2.10. For any integers n, k with $n \geq k \geq 0$, we have

$$B_{n,k}(B_1^{(L,Y)}(x), B_2^{(L,Y)}(x), \dots, B_{n-k+1}^{(L,Y)}(x)) = \sum_{j=k}^n \left\{ \begin{matrix} j \\ k \end{matrix} \right\}_{L_Y(n, j)} x^j.$$

From (22) and $k \geq 0$, we have

$$\begin{aligned} \sum_{n=0}^{\infty} \left(\frac{d}{dx} \right)^k B_n^{(L,Y)}(x) \frac{t^n}{n!} &= \left(\frac{d}{dx} \right)^k e^{x \left[E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right]} \\ &= k! \frac{1}{k!} \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)^k e^{x \left[E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right]} \\ &= k! \sum_{l=k}^{\infty} L_Y(l, k) \frac{t^l}{l!} \sum_{j=0}^{\infty} B_j^{(L,Y)}(x) \frac{t^j}{j!} = \sum_{n=k}^{\infty} \left(k! \sum_{j=0}^{n-k} B_j^{(L,Y)}(x) L_Y(n-j, k) \binom{n}{j} \right) \frac{t^n}{n!}. \end{aligned} \quad (32)$$

In particular, for $k = 1$, we get

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{d}{dx} B_n^{(L,Y)}(x) \frac{t^n}{n!} &= \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right) e^{x \left[E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right]} \\ &= \sum_{l=1}^{\infty} E[\langle Y \rangle_l] \frac{t^l}{l!} \sum_{j=0}^{\infty} B_j^{(L,Y)}(x) \frac{t^j}{j!} \\ &= \sum_{n=1}^{\infty} \left(\sum_{j=0}^{n-1} E[\langle Y \rangle_{n-j}] B_j^{(L,Y)}(x) \binom{n}{j} \right) \frac{t^n}{n!}. \end{aligned} \quad (33)$$

Therefore, by (32) and (33), we obtain the following theorem.

Theorem 2.11. For any integers n, k with $n \geq k \geq 0$, we have

$$\left(\frac{d}{dx} \right)^k B_n^{(L,Y)}(x) = k! \sum_{j=0}^{n-k} \binom{n}{j} B_j^{(L,Y)}(x) L_Y(n-j, k).$$

In particular, for $k = 1$, we get

$$\frac{d}{dx} B_n^{(L,Y)}(x) = \sum_{j=0}^{n-1} \binom{n}{j} E[\langle Y \rangle_{n-j}] B_j^{(L,Y)}(x), \quad (n \geq 1).$$

From (15), we have

$$\begin{aligned} \sum_{n=k}^{\infty} L_Y(n, k) \frac{t^n}{n!} &= \frac{1}{k!} \left(E \left[\left(\frac{1}{1-t} \right)^Y \right] - 1 \right)^k = \frac{1}{k!} \left(\sum_{j=1}^{\infty} E[\langle Y \rangle_j] \frac{t^j}{j!} \right)^k \\ &= \sum_{n=k}^{\infty} B_{n,k} \left(E[\langle Y \rangle_1], E[\langle Y \rangle_2], \dots, E[\langle Y \rangle_{n-k+1}] \right) \frac{t^n}{n!}. \end{aligned} \quad (34)$$

Let Y be the Poisson random variable with parameter $\alpha > 0$. Then we have

$$\begin{aligned} \sum_{n=0}^{\infty} E[\langle S_k \rangle_n] \frac{t^n}{n!} &= E\left[\left(\frac{1}{1-t}\right)^{S_k}\right] = E\left[\left(\frac{1}{1-t}\right)^{Y_1+\dots+Y_k}\right] \\ &= E\left[\left(\frac{1}{1-t}\right)^{Y_1}\right] E\left[\left(\frac{1}{1-t}\right)^{Y_2}\right] \cdots E\left[\left(\frac{1}{1-t}\right)^{Y_k}\right] \\ &= e^{-\alpha k} e^{\alpha k \left(\frac{1}{1-t}\right)} = e^{\alpha k \left(\frac{1}{1-t}-1\right)} = \sum_{n=0}^{\infty} B_n^L(\alpha k) \frac{t^n}{n!}. \end{aligned} \quad (35)$$

Therefore, by (35), we obtain the following theorem.

Theorem 2.12. Let Y be the Poisson random variable with parameter $\alpha > 0$. Then we have

$$E[\langle S_k \rangle_n] = B_n^L(\alpha k), \quad (n, k \geq 0).$$

From (22) and (26), we note that

$$\begin{aligned} \sum_{n=0}^{\infty} B_n^{(L,Y)}(x) \frac{t^n}{n!} &= e^{x \left(E\left[\left(\frac{1}{1-t}\right)^Y\right] - 1 \right)} = e^{x(e^{\alpha(\frac{1}{1-t}-1)} - 1)} \\ &= \sum_{k=0}^{\infty} \phi_k(x) \alpha^k \frac{1}{k!} \left(\frac{1}{1-t} - 1\right)^k = \sum_{k=0}^{\infty} \phi_k(x) \alpha^k \sum_{n=k}^{\infty} L(n, k) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \phi_k(x) \alpha^k L(n, k) \right) \frac{t^n}{n!}. \end{aligned} \quad (36)$$

Therefore, by (36), we obtain the following theorem.

Theorem 2.13. Let Y be the Poisson random variable with parameter $\alpha > 0$. For $n \geq 0$, we have

$$B_n^{(L,Y)}(x) = \sum_{k=0}^n \phi_k(x) \alpha^k L(n, k).$$

Assume that Y is the Bernoulli random variable with probability of success p . Then, we have

$$E[\langle Y \rangle_n] = \sum_{i=0}^1 \langle i \rangle_n p(i) = n!p, \quad (n \geq 1). \quad (37)$$

Thus, by (34) and (37), we get

$$\begin{aligned} L_Y(n, k) &= B_{n,k}(1!p, 2!p, \dots, (n-k+1)!p) \\ &= p^k B_{n,k}(1!, 2!, \dots, (n-k+1)!) = p^k L(n, k), \quad (n \geq k, n \geq 1), \end{aligned} \quad (38)$$

and, by (8) and (20), we have

$$B_n^{(L,Y)}(x) = \sum_{k=0}^n L_Y(n, k) x^k = \sum_{k=0}^n (px)^k L(n, k) = B_n^L(px), \quad (n \geq 1). \quad (39)$$

Therefore, by (39), we obtain the following theorem.

Theorem 2.14. Let Y be the Bernoulli random variable with probability of success p . Then we have

$$B_n^{(L,Y)}(x) = B_n^L(px), \quad (n \geq 1).$$

In addition, we have

$$L_Y(n, k) = p^k L(n, k), \quad (n \geq k, n \geq 1).$$

3. Conclusion

In this paper, by using generating functions we studied the probabilistic Lah numbers associated with Y and the probabilistic Lah-Bell polynomials associated with Y , as probabilistic versions of the Lah numbers and the Lah-Bell polynomials, respectively. Here Y is a random variable such that the moment generating function of Y exists in a neighborhood of the origin.

Our results indicate that the method of introducing random variables in [17, 18, 25] is applicable to various polynomials and numbers, and is also applicable to Lah numbers and Lah-Bell polynomials. The properties we derived, such as generating functions and recurrence relationships, are structurally very similar to the properties of probabilistic Bell polynomials([17]), probabilistic Dowling polynomials([25]) and probabilistic Jindalrae-Stirling polynomials([11]), highlighting the generality of this method.

As one of our future projects, we would like to continue to investigate degenerate versions, λ -analogues and probabilistic versions of many special polynomials and numbers and to find their applications to physics, science and engineering as well as to mathematics.

References

- [1] M. Abramowitz, I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series, No. 55, U. S. Government Printing Office, Washington, DC, 1964.
- [2] C. R. Adams, A. P. Morse, *Random sampling in the evaluation of a Lebesgue integral*, Bull. Amer. Math. Soc. **45** (1939), no. 6, 442–447.
- [3] J. A. Adell, *Probabilistic Stirling numbers of the second kind and applications*, J. Theoret. Probab. **35** (2022), no. 1, 636–652.
- [4] J. C. Ahuja, E. A. Enneking, *Concavity property and a recurrence relation for associated Lah numbers*, Fibonacci Q. **17** (1979), 158–161.
- [5] A. Aledamat, W. A. Khan, N. Ahmad, *Bell-based partially degenerate Genocchi polynomials and their applications*, Bull. Math. Anal. Appl. **16** (2024), no. 4, 61–76.
- [6] S. Araci, M. Acikgoz, *A note on the Frobenius-Euler numbers and polynomials associated with Bernstein polynomials*, Adv. Stud. Contemp. Math., Kyungshang **22** (2012), No. 3, 399–406.
- [7] K. Boubellouta, A. Boussayoud, S. Araci, M. Kerada, *Some theorems on generating functions and their applications*, Adv. Stud. Contemp. Math., Kyungshang **30** (2020), No. 3, 307–324.
- [8] K. N. Boyadzhiev, *Convolutions for Stirling numbers, Lah numbers, and binomial coefficients*, Proc. Jangjeon Math. Soc. **25** (2022), no. 2, 227–244.
- [9] L. Carlitz, *Some arithmetic properties of the Bell polynomials*, Bull. Amer. Math. Soc. **71** (1965), 143–144.
- [10] L. Comtet, *Advanced combinatorics*, Revised and enlarged edition, D. Reidel Publishing Co., Dordrecht, 1974.
- [11] W. A. Khan, U. Duran, N. Ahmad, *Probabilistic degenerate Jindalrae and Jindalrae-Stirling polynomials of the second kind*, J. Convex Nonlinear Anal. **26** (2025), no. 4, 941–961.
- [12] N. Kilar, Y. Simsek, *Combinatorial sums involving Fubini type numbers and other special numbers and polynomials: approach trigonometric functions and p -adic integrals*, Adv. Stud. Contemp. Math., Kyungshang **31** (2021), No. 1, 75–87.
- [13] D. S. Kim, H. K. Kim, T. Kim, H. Lee, S. Park, *Multi-Lah numbers and multi-Stirling numbers of the first kind*, Adv. Difference Equ. 2021 (2021), Paper No. 411.
- [14] D. S. Kim, T. Kim, *Lah-Bell numbers and polynomials*, Proc. Jangjeon Math. Soc. **23** (2020), No. 4, 577–586.
- [15] D. S. Kim, T. Kim, *r -extended Lah-Bell numbers and polynomials associated with r -Lah numbers*, Proc. Jangjeon Math. Soc. **24** (2021), No. 1, 1–10.
- [16] D. S. Kim, T. Kim, *Normal ordering associated with λ -Whitney numbers of the first kind in λ -shift algebra*, Russ. J. Math. Phys. **30** (2023), no. 3, 310–319.
- [17] D. S. Kim, T. Kim, *Probabilistic bivariate Bell polynomials*, Quaestiones Math. **48** (2025), no. 8, 1183–1196.
- [18] T. Kim, D. S. Kim, D. V. Dolgy, *Probabilistic degenerate derangement polynomials*, Math. Comput. Model. Dyn. Syst. **31** (2025), no. 1, article no. 2529188.
- [19] T. Kim, D. V. Dolgy, D. S. Kim, H. K. Kim, S. H. Park, *A note on degenerate generalized Laguerre polynomials and Lah numbers*, Adv. Difference Equ. 2021 (2021), Paper No. 421.
- [20] T. Kim, D. S. Kim, *Probabilistic degenerate Bell polynomials associated with random variables*, Russ. J. Math. Phys. **30** (2023), no. 4, (in press).
- [21] T. Kim, D. S. Kim, *Some identities involving degenerate Stirling numbers associated with several degenerate polynomials and numbers*, Russ. J. Math. Phys. **30** (2023), no. 1, 62–75.
- [22] T. Kim, D. S. Kim, D. V. Dolgy, J.-W. Park, *Degenerate binomial and Poisson random variables associated with degenerate Lah-Bell polynomials*, Open Math. **19** (2021), 1588–1597.
- [23] T. Kim, D. S. Kim, L.-C. Jang, H. Lee, H.-Y. Kim, *Complete and incomplete Bell polynomials associated with Lah-Bell numbers and polynomials*, Adv. Difference Equ. 2021 (2021), Paper No. 101.
- [24] Y. Ma, D. S. Kim, T. Kim, H. Kim, H. Lee, *Some identities of Lah-Bell polynomials*, Adv. Difference Equ. 2020 (2020), Paper No. 510.
- [25] Y. Ma, T. Kim, D. S. Kim, R. Xu, *Recurrence for probabilistic extension of Dowling polynomials*, Open Math. **23** (2025), no. 1, Paper No. 20250150, 7 pp.
- [26] S. Roman, *The umbral calculus*, Pure and Applied Mathematics, 111, Academic Press, Inc., New York, 1984.
- [27] S. M. Ross, *Introduction to probability models*, Eleventh edition, Elsevier/Academic Press, Amsterdam, 2014.

- [28] Y. Simsek, *Identities on the Changhee numbers and Apostol-type Daehee polynomials*, Adv. Stud. Contemp. Math., Kyungshang **27** (2017), No. 2, 199–212.
- [29] Y. Simsek, *Construction of generalized Leibnitz type numbers and their properties*, Adv. Stud. Contemp. Math., Kyungshang **31** (2021), No. 3, 311–323.
- [30] S. Tauber, *Lah numbers for R-polynomials*, Fibonacci Q. **6** (1968), No. 5, Sonderheft, 100–107.